

Strongly invertible knots, rational-fold branched coverings and hyperbolic spatial graphs

Kazuhiro ICHIHARA

School of Mathematics Education
Nara University of Education
Takabatake-cho
Nara 630-8528, JAPAN
ichihara@nara-edu.ac.jp

Akira USHIJIMA

Division of Mathematical and Physical Sciences
Graduate School of Natural Science and Technology
Kanazawa University
Kanazawa 920-1192, JAPAN
ushijima@kenroku.kanazawa-u.ac.jp

February 2, 2008

Abstract

A construction of a spatial graph from a strongly invertible knot was developed by the second author, and a necessary and sufficient condition for the given spatial graph to be hyperbolic was provided as well. The condition is improved in this paper. This enable us to show that certain classes of knots can yield hyperbolic spatial graphs via the construction.

Key words: spatial graph, theta-curve, strongly invertible knot, simple knot, hyperbolic manifold.

2000 Mathematics Subject Classifications: Primary: 57M50;
secondary: 05C10.

1 Introduction

Throughout this paper the category we will work with is differentiable. We start this section with recalling several basic terminologies frequently used in

this paper. Let M be a compact orientable three-dimensional manifold. If it has boundary, then we denote the boundary by ∂M . A *spatial graph* in M means the image of an embedding of a graph, which will be regarded as a one-dimensional cellular complex as usual. We often identify a spatial graph and the graph of its source in this paper; a vertex, an edge, etc. of a spatial graph G refers that of the graph appearing as the source of G . Spatial graphs are assumed to be finite throughout this paper.

A spatial graph is especially called a *link* if it is homeomorphic to a disjoint union of circles, and a link is particularly called a *knot* if its connected component is one. Let $N(G)$ be a regular neighborhood of a spatial graph G in M , and let $E(G)$ be the *exterior* of G in M , namely $E(G)$ is the closure of $M - N(G)$ in M . Suppose that any connected components of ∂M are closed orientable surfaces except spheres. Then a spatial graph G in M is said to be *hyperbolic* if $E(G)$ minus all the toral boundary components admits a complete hyperbolic structure of finite volume with geodesic boundary.

Let G be a spatial graph in the three-dimensional sphere S^3 , and φ an orientation-preserving self-diffeomorphism of the pair (S^3, G) . For a positive integer n , we denote by φ^n the map obtained by iterating φ n times. Then φ is said to be *periodic of order n* when φ^n is not the identity map on (S^3, G) for any integer m with $1 \leq m \leq n-1$, and φ^n is the identity map. It is known that if such φ has a fixed point, then the fixed point set is homeomorphic to a circle. See Chapter 1 of [MB84] for example. A spatial graph G is said to be *strongly periodic of order n* if there is a periodic self-diffeomorphism φ of order n such that the fixed point set of φ intersects with each connected component of G at exactly two points, and that n is the maximal period that (S^3, G) can admit with the same fixed point set. Such a self-diffeomorphism is called a *strong inversion* when $n = 2$, and strongly periodic links of order 2 are especially called *strongly invertible* links.

In [Ush04], the second author has developed a method, called *the $n/2$ -fold cyclic branched covering*, to construct strongly periodic spatial graphs in S^3 from a strongly invertible link L in S^3 . Let α be the axis of a strong inversion ι of the pair (S^3, L) . Then the construction consists of the following two steps:

Step 1. Take a quotient of (S^3, L) by ι . Then we have S^3/ι , which is homeomorphic to S^3 . In S^3/ι , we have a spatial graph G as the quotient of L by ι , which has two univalent vertices on $\alpha/\iota = \alpha$.

Step 2. Take the n -fold cyclic branched covering of $(S^3/\iota, G)$ along α . Then we have the n -fold cyclic branched covering of S^3/ι , which is again homeomorphic to S^3 . In the n -fold cyclic branched covering of S^3/ι , we have a spatial graph G' as the pre-image of G by the covering.

We call the spatial graph G' obtained by this procedure the *$n/2$ -fold cyclic branched covering of L along α* . A spatial graph is called a θ_n -*curve* for some integer $n \geq 3$ when each connected component has exactly two vertices of valency n . If we start with a strongly invertible link, then the $n/2$ -fold cyclic branched covering yields strongly periodic θ_n -curves. We have arcs as the $1/2$ -fold cyclic




Figure 1: The process of constructing $\frac{n}{2}$ -fold cyclic branched covering of the trefoil knot

branched covering when $n = 1$, and we then call the union of them as a strongly periodic θ_1 -curve in this paper. Similarly we have the original L as the $2/2$ -fold cyclic branched covering when $n = 2$, and we occasionally call it a strongly periodic θ_2 -curve.

2 When can we have strongly periodic hyperbolic θ_n -curves?

Let M be a three-dimensional manifold admitting a periodic self-diffeomorphism φ of order n . Then a submanifold S of M is said to be *equivariant under φ* if either $\varphi^i(S) = S$ or $\varphi^i(S) \cap S = \emptyset$ holds for any integer i with $1 \leq i \leq n - 1$. Let Γ be a finite group acting on M . Then S is said to be Γ -*equivariant* if it is equivariant under the action of any element of Γ .

Let K be a non-trivial strongly invertible knot in S^3 admitting a strong inversion with axis α , and θ_n^K the strongly periodic θ_n -curve obtained from K by the $n/2$ -fold cyclic branched covering along α . Then it has been essentially proved in [Ush04] that θ_n^K is hyperbolic for any $n \geq 3$ if and only if $E(K \cup \alpha)$ contains no essential tori which are equivariant under the action of the strong inversion.

2.1 Main theorem

The first purpose of this paper is to simplify and improve the result given in [Ush04]. A spatial graph in S^3 is said to be *trivial* when it can be ambient isotopic into an embedded two-dimensional sphere.

Theorem 1. *Let K be a non-trivial strongly invertible knot in S^3 admitting a strong inversion with axis α , and θ_n^K the strongly periodic θ_n -curve obtained from K by the $n/2$ -fold cyclic branched covering along α . Then the following conditions are equivalent:*

- (1) For any $n \geq 3$, θ_n^K is hyperbolic.
- (2) For some $n \geq 3$, θ_n^K is hyperbolic.
- (3) Any embedded torus in $E(K \cup \alpha)$ is compressible in $E(K \cup \alpha)$.
- (4) Any embedded equivariant torus in $E(K \cup \alpha)$ is compressible in $E(K \cup \alpha)$.
- (5) Any embedded torus in $E(K)$ which is disjoint from α is compressible in $E(K)$.
- (6) Any embedded equivariant torus in $E(K)$ which is disjoint from α is compressible in $E(K)$.

There are some remarks about this theorem.

- We shall use standard language in three-dimensional topology. We here include some of them for reader's convenience. See [Jac80] on such terminologies for example. Let M be a compact three-dimensional manifold.

A properly embedded surface F in M other than a sphere or a disc is said to be *compressible* in M if it admits a disc D such that $F \cap D = F \cap \partial D = \partial D$ and that ∂D is not null-homotopic on F . Such a disc is called a *compressing disc* for F . Otherwise F is said to be *incompressible*.

Analogously, a disc D' embedded in M is called a *boundary-compressing disk* for F if $\partial D' = a \cup b$, where $a := F \cap D' = F \cap \partial D'$ is an arc which is not parallel to ∂F in F and $b := \partial M \cap D' = \partial M \cap \partial D'$ is an arc in ∂M which is not homotopic into ∂F relative ∂b . If there is no boundary-compressing disk for F , then F is said to be *boundary-incompressible*.

The manifold M is said to be *irreducible* if any embedded sphere bounds an embedded ball, and *boundary-irreducible* if each connected component of ∂M is incompressible.

Two embedded surfaces F and F' are said to be *parallel* if they bound a product region in M . In particular, when F is parallel to a subsurface of ∂M , it is said to be *boundary-parallel*.

Finally, an embedded surface is said to be *essential* if it is incompressible, boundary-incompressible, and not boundary-parallel.

- Though we do not mention α in the notation θ_n^K , the spatial graph θ_n^K depends not only on both K and n but also on the choice of the axis α . See Subsection 3.2 for such an example.
- Since $\partial E(K \cup \alpha)$ consists of an orientable closed surface of genus three, there is no embedded torus which is boundary-parallel. Thus Condition (3) means that $E(K \cup \alpha)$ is *atoroidal*.

- The assumption that K is a *knot* is essential. Suppose that K is a two-component non-trivial torus link and α is the axis of the strong inversion. See Figure 2 for example. Then K is known to be strongly invertible, and $E(K \cup \alpha)$ is known to be atoroidal. But it has been proved as Theorem 1.3 in [Ush04] that θ_n^K is not hyperbolic.




Figure 2: Two-component torus link and its axis of a strong inversion

2.2 Equivariant Torus Theorem

In this subsection we give our version of the so-called “Equivariant Torus Theorem”. It should be emphasized that the boundary of the manifold in the proposition below is not assumed to be incompressible.

Proposition 2. *For a compact orientable irreducible three-dimensional manifold with non-empty boundary, if it admits an action of a finite group and if it contains an essential torus, then it contains an embedded equivariant incompressible torus as well.*

To prove this proposition, we prepare a definition. Let M be an orientable three-dimensional manifold, and let S be an essential torus properly embedded in M . Then we say that S is said to be *canonical* if any other properly embedded essential torus or annulus can be isotoped to be disjoint from S .

Lemma 3. *For a compact orientable irreducible three-dimensional manifold, if it admits an action of a finite group and if it contains a canonical torus, then the torus is isotoped to be equivariant under the action.*

Proof. Let M be a compact orientable irreducible three-dimensional manifold admitting an action of a finite group Γ , and T a canonical torus in M . Then we can choose a Γ -invariant Riemannian metric on M whose boundary has non-negative mean curvature. Such a metric can be constructed by averaging any metric over the action. Then we have a least area torus T' isotopic to T by virtue of the result in [FHS83]. Precisely we use an extension of Theorems 1.1 and 1.2 of [FHS83] to the non-empty boundary case, which is explained at the

3rd paragraph in page 635 of [FHS83]. Since the torus T' is also canonical as is T , its images by the action are isotopically disjoint or coincident, and thus they are actually distinct or coincident by an extension of Theorem 6.2 of [FHS83] to the non-empty boundary case, which is also explained at the 3rd paragraph in page 635 of [FHS83]. This indicates that T' is equivariant under the action. \square

Proof of Proposition 2. Let M be a compact orientable irreducible three-dimensional manifold admitting an action of a finite group, and T an essential torus in M .

If M contains a canonical torus, then it is isotoped to be equivariant under the action by Lemma 3. So we assume that M has no canonical torus from now on.

Let us set $M' := M$ if ∂M is incompressible. If, on the other hand, ∂M is compressible, we set M' as follows. We consider a maximal set of compression discs for ∂M , and compress ∂M along the discs. Then we have a sub-manifold N , possibly disconnected, in M . By standard cut-and-paste argument together with the assumption of the irreducibility for M , any incompressible surface in M can be isotoped apart from ∂N . So we can find a torus T' in N which is isotopic to T in M . Then we set M' as the connected component of N containing T' .

Here we note that any toral component of $\partial M'$ is boundary-parallel in M . For, as mentioned above, any incompressible surface in M can be isotoped apart from $\partial M' \subset \partial N$. So the toral component must be boundary-parallel in M by the assumptions that there is no canonical torus in M and that $\partial M'$ is incompressible in M by its construction. This implies that T' is essential in M' since T' is not boundary-parallel in M as is T .

As a result we thus find an irreducible three-dimensional submanifold M' containing an essential torus T' but not containing any canonical torus. Then M' must be either a Seifert fibred space or an I -bundle by Proposition 3.2 of [NS97]. Thus all connected components of $\partial M'$ are tori and all of them are, as claimed above, boundary-parallel in M . This concludes that $M' = M$, and thus M contains an incompressible torus as a boundary component, which gives an embedded equivariant incompressible torus. This completes the proof of Proposition 2. \square

2.3 Lemmata for the proof of Theorem 1

In this subsection, we will prepare three lemmata to prove Theorem 1.

The first one would be well-known in three-dimensional manifold theory. Actually a similar result is shown as Theorem in [GL84].

Lemma 4. *Suppose that an irreducible three-dimensional manifold admitting an action of a finite cyclic group contains an embedded torus equivariant under the action. If the torus is compressible, then there exists a compression disc which is equivariant under the action.*

Proof. Let M be an irreducible three-dimensional manifold admitting a non-trivial action φ of period n , and let T be an embedded torus in M which is equivariant under φ with compression disc D .

We first suppose that D has non-empty intersection with another $\varphi^i(T)$ for some $i \in \{1, 2, \dots, n-1\}$. Since $\varphi^i(T)$ are mutually disjoint, the intersection appearing on D must consist of circles in the interior of D . Then, since M is irreducible, one can reduce the circles which are inessential on the torus by isotopy. Take an inner-most one on D , and replace T and D with $\varphi^i(T)$ and the chosen disc. Thus, without loss of generality, we can assume that there is a compression disc for such a torus having no intersection with any other tori $\varphi^i(T)$, and we denote a pair of the torus and the compression disc by T and D again.

Take an equivariant regular neighborhood $N(T)$ of T under φ , and we define a submanifold W in M as follows:

$$W := M - \bigcup_{i=1}^n \varphi^i(N(T)).$$

Here we take $N(T)$ sufficiently small so that $D \cap W$ gives a compression disc in W for $\partial_0 W$, where $\partial_0 W$ is a connected component of ∂W given by one of two connected components of $\partial N(T)$. We then apply the Equivariant Loop Theorem (see [YM84]) to $(W, \partial W, D \cap W)$ so that we can find a new compression disc D' for $\partial_0 W$ in W , which is equivariant under φ . Enlarging this disc and we can find a compression disc for T in M , which is equivariant under φ . \square

We can prove the following two lemmata using this lemma.

Lemma 5. *Let L be a strongly invertible link in S^3 admitting a strong inversion with axis α , and θ_n^L the spatial graph with axis α_n which is obtained from L by $n/2$ -fold cyclic branched covering along α . Then, if any embedded equivariant torus in $E(\theta_n^L \cup \alpha_n)$ is compressible in $E(\theta_n^L \cup \alpha_n)$ for some $n \geq 1$, then so is it for all $n \geq 1$.*

Remark. Strongly periodic action on $E(\theta_1^L \cup \alpha_1)$ means the trivial action. So an embedded *equivariant* torus in $E(\theta_1^L \cup \alpha_1)$ is nothing but an embedded torus, or in other words, any embedded torus is an equivariant torus in $E(\theta_1^L \cup \alpha_1)$ in our sense.

Proof. Our proof consists of the following two steps:

Step 1. Suppose that any embedded equivariant torus in $E(\theta_n^L \cup \alpha_n)$ is compressible for some $n \geq 2$, and prove that so is it in $E(\theta_1^L \cup \alpha_1)$.

Step 2. Suppose that any embedded equivariant torus in $E(\theta_1^L \cup \alpha_1)$ is compressible, and prove that so is it in $E(\theta_n^L \cup \alpha_n)$ for any $n \geq 2$.

By taking a regular neighborhood suitably, we can assume that $E(\theta_n^L \cup \alpha_n)$ is an n -fold cyclic covering space of $E(\theta_1^L \cup \alpha_1)$.

To prove the first step, let T be an embedded (equivariant) torus in $E(\theta_1^L \cup \alpha_1)$, and we will find its compression disc in $E(\theta_1^L \cup \alpha_1)$. Consider the preimage of T in $E(\theta_n^L \cup \alpha_n)$ by the covering projection. It consists of either an equivariant

torus, or a tuple of mutually disjoint equivariant tori. Let \tilde{T} be one of its connected components. Then the assumption says that there is a compression disc for \tilde{T} in $E(\theta_n^L \cup \alpha_n)$. Furthermore $E(\theta_n^L \cup \alpha_n)$ is irreducible since so is S^3 and $E(\theta_n^L \cup \alpha_n) \subset S^3$. So we can apply Lemma 4 and find an equivariant compression disc for \tilde{T} in $E(\theta_n^L \cup \alpha_n)$, and the projection of the disc in $E(\theta_1^L \cup \alpha_1)$ becomes the compression disc for T in $E(\theta_1^L \cup \alpha_1)$. We have thus proved the first step.

To prove the second step, let T be an embedded equivariant torus in $E(\theta_n^L \cup \alpha_n)$, and we will find its compression disc in $E(\theta_n^L \cup \alpha_n)$. Since T has no intersection with the axis α_n of φ , the projection of T by the quotient map induced by the action of φ is an embedded torus in $E(\theta_1^L \cup \alpha_1)$. Since we are assuming that any embedded torus in $E(\theta_1^L \cup \alpha_1)$ is compressible, there is a compression disc for the projection of T . Since the disc has no intersection with the branched set α_1 , its lift in $E(\theta_n^L \cup \alpha_n)$ is a compression disc for T . We have thus proved the second step, which means that the proof of Lemma 5 have been completed. \square

Lemma 6. *Let G be a strongly periodic spatial graph in S^3 admitting a periodic action with axis α . If an embedded equivariant torus in $E(G)$ which is disjoint from α is compressible in $E(G)$, then it has a compression disc which is disjoint from α as well.*

Proof. Let φ be the periodic of order n on (S^3, G) with axis α . By taking a regular neighborhood of G suitably, we can assume that $E(G)$ admits the action of φ . Let T be an embedded torus in $E(G)$ which is equivariant under φ . Now we have its compression disc in $E(G)$ by the assumption of this lemma. Since $E(G)$ is irreducible as is S^3 , we can find an equivariant compression disc D for T by Lemma 4.

If D has no intersection with α , then we are done. So we assume $D \cap \alpha \neq \emptyset$ from now on. Since α is the fixed point set of the action on (S^3, G) and since D is equivariant under this action, we can say that $\varphi^i(D) = D$ holds for any $i \in \{1, 2, \dots, n\}$, i.e., $\varphi|_D$ can be regarded as a periodic diffeomorphism on D . Then, by the well-known classical result, there is an element r in the orthogonal group of dimension two together with a homeomorphism h on D such that $\varphi|_D = h r h^{-1}$. See Theorem 3.1 in [CK94] for its modern proof. So we have the following two cases, depending on the orientation of $\varphi|_D$ on D :

- $\varphi|_D$, and thus r , is orientation-reversing, or
- $\varphi|_D$, and thus r , is orientation-preserving.

If the first case happens, the fixed point set of r is a diameter of D and thus the fixed point set of $\varphi|_D$ is an embedded arc on D , which is $D \cap \alpha$. We take a sufficiently small regular neighborhood $N(\alpha)$ in $E(G)$ so that $D \cap \partial N(\alpha)$ consists of two arcs parallel to $D \cap \alpha$. Then replace $D \cap N(\alpha)$ by one of two simply-connected regions on $\partial N(\alpha)$ bounded by four arcs: two of them come from $T \cap (\partial N(\alpha) - D)$ and the other two arcs are parallel ones appearing as


$D \cap \partial N(\alpha)$. We thus have a compression disc for T which has no intersection with α .

We lastly consider the second case. Since r is non-trivial, the set of fixed points for r consists of a single point, and so does the set of fixed points for $\varphi|_D$. This means that $D \cap \alpha$ consists of a single point. The axis α is the unknot by the affirmative solution of the Smith Conjecture. See page 4 of [MB84]. So we can regard T as the boundary of a trivially-embedded solid torus V with core curve α . Since $G \cup \alpha$ is connected by the definition of strongly periodic θ_n -curve, and since $T \cap \alpha = \emptyset$, V entirely contains $G \cup \alpha$. So the closure of $E(G \cup \alpha) - V$ is also a trivially-embedded solid torus. Therefore we can find a compression disc for $T = \partial V$ in $E(G \cup \alpha) - V \subset E(G \cup \alpha)$, which induces a compression disc for T in $E(G \cup \alpha)$. \square

2.4 Proof of Theorem 1

Now we are in the position to give a proof of Theorem 1.

Proof of Theorem 1. Though the equivalence between (1) and (4) has been partially proved as Theorem 1.2 in [Ush04], we will give a whole proof of Theorem 1 for completeness. We show it following the diagram below:



(1) \rightarrow (2). This is clear.

(2) \rightarrow (4). Assume that the strongly periodic θ_n -curve θ_n^K is hyperbolic for some $n \geq 3$. Then there is no essential torus in $E(\theta_n^K)$ by Thurston's study of geometric structure on three-dimensional manifolds. So any embedded equivariant torus in $E(\theta_n^K)$ which is disjoint from α_n is compressible in $E(\theta_n^K)$ as well. Then there is a compression disc in $E(\theta_n^K)$ which has no intersection with α_n by Lemma 6. This means that any embedded equivariant torus in $E(\theta_n^K \cup \alpha_n)$ is compressible in $E(\theta_n^K \cup \alpha_n)$. So we can apply Lemma 5 for $n = 2$ and we thus have (4) since $E(\theta_2^K \cup \alpha_2) = E(K \cup \alpha)$.

(4) \rightarrow (1). Since θ_1^K is an arc whose terminal vertices lie on α_1 , the manifold $E(\theta_1^K)$ is topologically the three-dimensional ball, and α_1 turns out to be two arcs in the ball, which we will denote by α_1 again. Thus $(E(\theta_1^K), \alpha_1)$ can be regarded as a two-string tangle. This realization is the so-called Montesinos trick. See [Mon75, BH96].

Since K is a non-trivial knot in S^3 , the exterior of K in S^3 is irreducible and boundary-irreducible. The two-string tangle $(E(\theta_1^K), \alpha_1)$ is thus prime by Theorem 3.5.17 in [Kaw96].

Moreover Lemma 5 can be applied by the assumption so that any embedded equivariant torus in $E(\theta_1^K \cup \alpha_1)$ is compressible in $E(\theta_1^K \cup \alpha_1)$. Since the finite group acting on $E(\theta_1^K \cup \alpha_1)$ is the trivial group, the result above actually means that any embedded torus in $E(\theta_1^K \cup \alpha_1)$ is compressible in $E(\theta_1^K \cup \alpha_1)$. This implies that the tangle $(E(\theta_1^K), \alpha_1)$ atoroidal, or it is the so-called *hyperbolic* tangle in the sense of [Kaw96].

Let $(D_{E(\theta_1^K)}, D_{\alpha_1})$ be the double of $(E(\theta_1^K), \alpha_1)$ along the sphere $\partial E(\theta_1^K)$. Then D_{α_1} gives a hyperbolic link in S^3 by Exercise 3.6.4 together with Theorem 3.6.6 both in [Kaw96]. Now, as we saw in Theorem 2.1 in [Ush04], the n -fold cyclic branched covering of $(D_{E(\theta_1^K)}, D_{\alpha_1})$ along D_{α_1} is a closed hyperbolic manifold for any $n \geq 3$.

Each manifold obtained by the covering admits an involution along the surface in the manifold induced from $\partial E(\theta_1^K)$. The action can be regarded as an isometry by Mostow-Prasad's rigidity theorem (see [Fri04] for example). Thus the quotient by this action is a hyperbolic manifold with totally geodesic boundary. As a result, the n -fold cyclic branched covering of $(E(\theta_1^K), \alpha_1)$ along α_1 is a hyperbolic manifold with totally geodesic boundary for any $n \geq 3$. This conclusion means that (1) holds by the definition of $n/2$ -fold cyclic branched covering.

- (3) \rightarrow (5). This is trivial since $E(K \cup \alpha)$ is a submanifold in $E(K)$.
- (5) \rightarrow (6). This is trivial.
- (6) \rightarrow (4). This is an immediate consequence of Lemma 6.
- (4) \rightarrow (3). Proposition 2 guarantees the contraposition of this statement as follows: suppose that there is an incompressible torus in $E(K \cup \alpha)$. Since the genus of $\partial E(K \cup \alpha)$ is three, it is not boundary-parallel, i.e., it is actually essential. Then we can find an embedded equivariant incompressible torus in $E(K \cup \alpha)$ by Proposition 2.

We have thus completed the proof of Theorem 1. \square

Let G_n be a hyperbolic strongly periodic θ_n -curve in S^3 . Then, as in the previous theorem, a knot can be obtained from G_n by the $2/n$ -fold cyclic branched covering; namely first take the quotient by the periodic diffeomorphism of order n , and then take the double branched covering along the axis of the periodic diffeomorphism. The knot obtained by this procedure must be a non-trivial strongly invertible knot, and the spatial graph obtained from this knot by $l/2$ -fold cyclic branched covering is a hyperbolic strongly periodic θ_l -curve, which can be called the *spatial periodic θ_l -curve obtained from G_n by the l/n -fold cyclic branched covering*. We have thus obtained the following corollary as an immediate consequence of Theorem 1:

Corollary 7. *Let G_n be a hyperbolic strongly periodic θ_n -curve in S^3 . Then, for any integer $l \geq 3$, G_l is a hyperbolic strongly periodic θ_l -curve in S^3 as well, where G_l is constructed from G_n by the l/n -fold cyclic branched covering. \square*

Now, once we have obtained Theorem 1, the following question naturally arises:

Question. Which strongly invertible knot satisfies the conditions of Theorem 1?

We will give several partial answers to this question in the next section.

3 Which strongly invertible knot can be a source of strongly periodic hyperbolic θ_n -curves?

We start this section with proving the following lemma.

Lemma 8. *Let G be a strongly periodic spatial graph in S^3 admitting a periodic action with axis α . Suppose that there is an embedded equivariant torus in $E(G)$ which is disjoint from α and boundary-parallel. Then G must be a knot, and the torus has a compression disc in $E(G)$; this implies that G is the trivial knot.*

Proof. The assumption of this lemma says that there is an embedded torus, say T , in $E(G \cup \alpha)$ which is parallel to a boundary component $\partial_0 E(G)$ of $\partial E(G)$ in $E(G)$. This means that G has at least one connected component which is homeomorphic to a circle. Thus, by the definition of strongly periodic spatial graph, the strongly periodic action must be a strong inversion, which we denote by ι . We now start this proof with showing that G is actually a knot.

By taking regular neighborhoods of G and α suitably, we can assume that both $E(G)$ and $E(G \cup \alpha)$ admit an action induced by the periodic diffeomorphism. Since T is parallel to $\partial_0 E(G)$, there is a compact submanifold V in $E(G)$ such that ∂V consists of T and $\partial_0 E(G)$ and that V is homeomorphic to the product manifold $T^2 \times [0, 1]$ constructed by the two-dimensional torus T^2 . Since T can be regarded as in S^3 , it separates S^3 into two compact submanifolds. We denote one of them containing V by M_V . Then, since $\partial_0 E(G) \cap \alpha \neq \emptyset$ and $T \cap \alpha = \emptyset$ whereas $G \cup \alpha$ is connected, M_V entirely contains $G \cup \alpha$. Thus all boundary components of $E(G)$ must be contained in M_V . However, since M_V consist of the union of a solid torus and V , M_V can contain at most one component of G . This implies that G must be a knot, and thus $\partial_0 E(G) = \partial E(G)$.

We next show that the torus T has a compression disc in $E(G)$. Since T is equivariant under ι , possible arrangements between T and $\iota(T)$ are either $\iota(T) = T$ or $\iota(T) \cap T = \emptyset$. So one of the following two cases can happen for $\iota(T)$:

- $\iota(T) \subset V$, or
- $\iota(T) \subset E(G) - V$.

We first consider the latter case. Then, since ι is a homeomorphism which preserves $\partial E(G)$, the submanifold $\iota(V)$ is again homeomorphic to $T^2 \times [0, 1]$ with $\partial \iota(V) = \iota(T) \cup \partial E(G)$. Thus $\iota(T)$ is again parallel to $\partial E(G)$. Furthermore, since $\iota = \iota^{-1}$ and $\iota(T) \cap T = \emptyset$ in this case, the embedded torus T must be contained

in $\iota(V)$. So we can regard $\iota(V)$ and $\iota(T)$ as V and T , and the situation turns out to be the former case.

We then consider the former case. Recall that what we want to prove is the existence of a compression disc for T in $E(G)$, and we will do it by contradiction. So we assume that T is incompressible in $E(G)$. Then, since ι is a homeomorphism, $\iota(T)$ is also incompressible in $E(G)$, and thus especially incompressible in V . Since V has a homeomorphism ψ to $T^2 \times [0, 1]$, the embedded torus $\psi \circ \iota(T)$ is also incompressible in $T^2 \times [0, 1]$. Then, by LEMMA (1) in [Hei81], any incompressible surface in $T^2 \times [0, 1]$ must be parallel to both $T^2 \times \{0\}$ and $T^2 \times \{1\}$, and thus so is $\psi \circ \iota(T)$. Since $\partial E(G)$ is preserved by ι , we have $\psi \circ \iota(V) \subseteq T^2 \times [0, 1]$, which is equivalent to $\iota(V) \subseteq \psi^{-1}(T^2 \times [0, 1]) = V$. Thus we have the following relation:

$$V = \iota^2(V) \subseteq \iota(V) \subseteq V,$$

where we used the fact that ι^2 is trivial.

So we have $\iota(V) = V$, which means that $\iota|_V$ is an order two periodic action on V with $\iota(\partial E(G)) = \partial E(G)$ and $\iota(T) = T$. Thus we have $\partial(T^2 \times [0, 1]) = \psi(\partial E(G)) \cup \psi(T)$ and $\psi \circ \iota \circ \psi^{-1}$ is an order two periodic action on $T^2 \times [0, 1]$ which preserves both $\psi(\partial E(G))$ and $\psi(T)$. Then it is shown as Theorem 2.1 in [MS86] that any finite group action on a three-dimensional manifold with product structure preserves the structure. So $\psi(T)$ must intersect with the axis $\psi(V \cap \alpha)$ of the order two periodic action. This is equivalent to $T \cap \alpha \neq \emptyset$, which contradicts the assumption that $T \subset E(G \cup \alpha)$. Thus T is compressible in $E(G)$.

We have thus proved Lemma 8 □

3.1 Simple knots and tunnel number one knots

A three-dimensional manifold is said to be *simple* when it contains no essential torus. A link is said to be *simple* when it is non-trivial and its exterior is a simple manifold. By Thurston's study of geometric structures on knot exterior in S^3 , any non-trivial knot in S^3 is known to be one of a torus knot, a hyperbolic knot or a satellite knot. See [Thu82]; see also [Ota01, Kap01] for a detailed proof. Thus any simple knot in S^3 is either a torus knot or a hyperbolic knot. For such a knot, we have the following proposition:

Proposition 9. *For any strongly invertible non-trivial simple link L in S^3 and the axis α of any strong inversion of the pair (S^3, L) , any embedded equivariant torus in $E(L)$ which is disjoint from α is compressible in $E(L)$.*

Proof. The definition of simpleness for a link L implies that there is no essential torus in $E(L)$. So any embedded equivariant torus in $E(L)$ is either compressible or boundary-parallel. However, since L is non-trivial, the latter case cannot happen by Lemma 8. □

A knot K in S^3 is said to be *tunnel number one* if there exists an arc τ properly embedded in $S^3 - K$ such that $E(K \cup \tau)$ becomes a handlebody of

genus two. Such a knot is isotopic onto the genus two surface appearing as the boundary $\partial E(K \cup \tau)$, which gives a standard genus two Heegaard surface in S^3 . So the knot admits a strong inversion induced from that of the surface. Thus every tunnel number one knot is known to be strongly invertible. See also Lemma 5 in [Mor95].

Proposition 10. *For any non-trivial tunnel number one knot K in S^3 and the axis α of any strong inversion of the pair (S^3, K) , any embedded equivariant torus in $E(K)$ which is disjoint from α is compressible in $E(K)$.*

Proof. Because of Proposition 9, we only consider the case where the tunnel number one knot is satellite. For such a knot K , it has been proved as Theorem 2.1 in [MS91] that it is made of a non-trivial torus knot K_0 and a two-bridge link $K_1 \cup K_2$ which is neither a trivial link nor a Hopf link by gluing $\partial E(K_2)$ to $\partial E(K_0)$. Since K_2 itself is a trivial knot in S^3 , $E(K_2)$ is a solid torus containing K_1 . Thus the image of K_1 under this gluing can be regarded as a knot in S^3 . From now on, we will denote the images of $E(K_1 \cup K_2)$ and $E(K_2)$ under this gluing by the same symbols. Then we say that $E(K)$ is decomposed into two parts; one is the torus knot exterior part $E(K_0)$, which is a Seifert fibered space, and the other is the two-bridge link exterior part $E(K_1 \cup K_2)$, which admits hyperbolic structure. Thus the characteristic submanifold of $E(K)$ coincides with $E(K_0)$ up to ambient isotopy in $E(K)$. See [Jac80]. Then, by Theorem 8.6 in [MS86], $E(K_0)$ can be taken to be preserved by the strong inversion ι of the pair (S^3, K) with axis α . This implies that the action of ι gives a periodic action of order two on $E(K_0)$, and it then equals to the restriction to $E(K_0)$ of the strong inversion of the torus knot. See 10.6 of [Kaw96] for example.

Let T be an embedded equivariant torus in $E(K)$ which is disjoint from α . If T is not compressible in $E(K)$, then one of the following two cases can happen:

- The torus T is essential in $E(K)$, or
- The torus T is boundary-parallel in $E(K)$.

If the latter case happens, then we can apply Lemma 8 so that K must be a trivial knot. This contradicts the assumption of K . So we study the former case from now on. Then this assumption implies that T is essential in $E(K \cup \alpha)$ as well. Since the satellite torus T_s is incompressible in $E(K)$, the punctured surface $T'_s := T_s \cap E(K \cup \alpha)$ is also incompressible in $E(K \cup \alpha)$. Thus, in the former case, T can be isotoped in $E(K \cup \alpha)$ so that one of the following two subcases occurs:

- The torus can be isotoped to T' so that the two tori T' and T_s intersect each other, and the intersection consists of circles essential in both T' and T'_s .
- The torus can be isotoped to T' so that the two tori T' and T_s do not intersect each other.

We start with considering the former subcase. Then we can also assume that the intersection is minimal up to isotopy in $E(K \cup \alpha)$. Then $T' \cap E(K_0)$ consists of essential annuli in $E(K_0) - (E(K_0) \cap \alpha)$. By taking the projection of $(E(K_0), \alpha)$ by the quotient map induced from ι , we have a two-string tangle (B^3, t) .

Claim. This tangle (B^3, t) is prime and atoroidal.

Proof. Since $E(K_0)$ is irreducible and boundary-irreducible, (B^3, t) is prime. See Theorem 3.5.15 in [Kaw96] for example. Since torus knots are simple knots, $E(K_0)$ is atoroidal. Thus any embedded equivariant torus in $E(K_0)$ which is disjoint from α is either boundary-parallel or compressible in $E(K_0)$. If the former case happens, then K_0 must be the trivial knot by Lemma 8. So such a torus is always compressible in $E(K_0)$. Furthermore, we can always find such a compression disc in $E(K_0)$ with no intersection with α by Lemma 6. This means that the exterior of $E(K_0) \cap \alpha$ in $E(K_0)$ is also atoroidal. Then Lemma 5 can be applied so that any embedded torus in (B^3, t) is compressible in (B^3, t) , which means that (B^3, t) is atoroidal. \square

On the other hand, each essential annulus appearing as a component of $T' \cap E(K_0)$ descends into $B^3 - t$ as an immersed essential, i.e., π_1 -injective, annulus. Then we can find an embedded essential annulus in $B^3 - t$ by the Annulus Theorem (see VIII.13 of [Jac80] for example). However this contradicts the following fact: if (B^3, t) is prime and atoroidal, then $B^3 - t$ contains no essential annulus. See 3.6.4 of [Kaw96] for example.

Thus the former subcase cannot occur and now we assume the latter subcase, or equivalently, T is isotoped into $E(K_0 \cup \alpha)$ or $E(K_1 \cup K_2 \cup \alpha)$. Recall that the restriction of the strong inversion ι into $E(K_0)$ (resp. $E(K_1 \cup K_2)$) gives its strong inversions with axis $\alpha \cap E(K_0)$ (resp. $\alpha \cap E(K_1 \cup K_2)$). Then, since T is essential, we can apply Proposition 2 so that there is an equivariant essential torus. This contradicts Proposition 9 since non-trivial torus knots are known to be simple; actually they are *small* (see Subsection 3.2), and non-trivial two-bridge links except Hopf link are also known to be simple (see Corollary 5 of [Oer84] for example).

We have thus proved Proposition 10. \square

These two propositions together with Theorem 1 (6) immediately implies the following result:

Theorem 11. *Let K be a non-trivial and strongly invertible knot in S^3 . If K is simple or tunnel number one, then, for any $n \geq 3$, the strongly periodic θ_n -curve constructed by the $n/2$ -fold cyclic branched covering along any axis of a strong inversion of the pair (S^3, K) is hyperbolic.* \square

As a corollary to Propositions 9 and 10, we have the following result, which gives a necessary condition for a non-trivial strongly invertible knot to be simple or tunnel number one.

Corollary 12. *Let K be a non-trivial strongly invertible knot in S^3 admitting a strong inversion ι with axis α . Let G be the spatial θ_3 -curve appearing as the quotient of $K \cup \alpha$ under ι . Then, if K is simple or tunnel number one, any embedded torus in $E(G)$ is compressible.*

Proof. Take an embedded torus T in $E(G)$, and let \tilde{T} be a torus in $E(K \cup \alpha)$ appearing as a component of the pre-image of T by ι . The torus \tilde{T} is equivariant under the action of the involution ι , and disjoint from α in $E(K)$ by construction. Since K is assumed to be simple or tunnel number one, we can find a compression disc for \tilde{T} which is disjoint from α by Propositions 9 and 10 and Theorem 1. Then \tilde{T} has a compression disc which is disjoint from α and equivariant under the action of ι by Lemma 4, which descends to a compression disc for T . \square

3.2 How about other knots?

Since we have Theorem 11, remaining strongly invertible knots are those which are satellite with tunnel number greater than one. In this case the property of hyperbolicity of the spatial graphs generally depends on the choice of the strong inversion.

One of its typical example is a *Granny knot*. This knot is made from a trefoil knot and its copy by their connected sum; see Figure 3. A knot in S^3 is said to be *small* if its exterior contains no closed essential surface. Since non-trivial torus knots are shown to be small by Theorem in [Tsa94], so is the trefoil knot. Furthermore, since it is tunnel number one, we can apply Theorem 4 in [MS00] so that the Granny knot is not tunnel number one.

The Granny knot K has two distinct strong inversion; one is with axis α , and the other is with axis β in Figure 3. If we choose the strong inversion with axis α , then we can see that the spatial graphs obtained by $n/2$ -fold cyclic branched covering are hyperbolic, by checking that the link D_{α_1} , which appeared in the proof of Theorem 1, is hyperbolic.

On the other hand, for $E(K \cup \beta)$, one can easily see that there is an essential torus, which is obtained from the decomposing sphere by tubing along one of the trefoil knot. This torus is the so-called *swallow-follow torus*. Thus we cannot obtain hyperbolic spatial graphs from this strong inversion by Theorem 1.

Thus we can say that the hyperbolicity depends on the choice of the strong inversion in general.

Acknowledgements

The first author is partially supported by Grant-in-Aid for Young Scientists (B), No. 18740038, Ministry of Education, Culture, Sports, Science and Technology, Japan.

The authors would like to thank Professor Kimihiko Motegi for his encouragement and comments, especially for his comments on Lemma 6, thank Professor Makoto Sakuma for his encouragement and comments, especially for Propo-




Figure 3: Connected sum of two trefoil knots with two distinct axes α and β of distinct strong inversions

sition 10, and thank Professor Ryo Nikkuni for his comments as a specialist of the spatial graph theory.

The authors also would like to thank the referee for his/her comments and suggestions, especially for the comment on the proof of Lemma 6.

References

- [BH96] Steven A. Bleiler and Craig D. Hodgson, *Spherical space forms and Dehn filling*, *Topology* **35** (1996), no. 3, 809–833.
- [CK94] Adrian Constantin and Boris Kolev, *The theorem of Kerékjártó on periodic homeomorphisms of the disc and the sphere*, *Enseign. Math.* (2) **40** (1994), no. 3-4, 193–204.
- [FHS83] Michael Freedman, Joel Hass, and Peter Scott, *Least area incompressible surfaces in 3-manifolds*, *Invent. Math.* **71** (1983), no. 3, 609–642.
- [Fri04] Roberto Frigerio, *Hyperbolic manifolds with geodesic boundary which are determined by their fundamental group*, *Topology Appl.* **145** (2004), no. 1-3, 69–81.
- [GL84] C. McA. Gordon and R. A. Litherland, *Incompressible surfaces in branched coverings*, *The Smith conjecture* (New York, 1979), *Pure Appl. Math.*, vol. 112, Academic Press, Orlando, FL, 1984, pp. 139–152.
- [Hei81] W. Heil, *Normalizers of incompressible surfaces in 3-manifolds*, *Glas. Mat. Ser. III* **16(36)** (1981), no. 1, 145–150.
- [Jac80] William Jaco, *Lectures on three-manifold topology*, *CBMS Regional Conference Series in Mathematics*, vol. 43, American Mathematical Society, Providence, R.I., 1980.

- [Kap01] Michael Kapovich, *Hyperbolic manifolds and discrete groups*, Progress in Mathematics, vol. 183, Birkhäuser Boston Inc., Boston, MA, 2001.
- [Kaw96] Akio Kawauchi, *A survey of knot theory*, Birkhäuser Verlag, Basel, 1996, Translated and revised from the 1990 Japanese original by the author.
- [MB84] John W. Morgan and Hyman Bass (eds.), *The Smith conjecture*, Pure and Applied Mathematics, vol. 112, Academic Press Inc., Orlando, FL, 1984, Papers presented at the symposium held at Columbia University, New York, 1979.
- [Mon75] José M. Montesinos, *Surgery on links and double branched covers of S^3* , Knots, groups, and 3-manifolds (Papers dedicated to the memory of R. H. Fox), Princeton Univ. Press, Princeton, N.J., 1975, pp. 227–259. Ann. of Math. Studies, No. 84.
- [Mor95] Kanji Morimoto, *There are knots whose tunnel numbers go down under connected sum*, Proc. Amer. Math. Soc. **123** (1995), no. 11, 3527–3532.
- [MS86] William H. Meeks, III and Peter Scott, *Finite group actions on 3-manifolds*, Invent. Math. **86** (1986), no. 2, 287–346.
- [MS91] Kanji Morimoto and Makoto Sakuma, *On unknotting tunnels for knots*, Math. Ann. **289** (1991), no. 1, 143–167.
- [MS00] Kanji Morimoto and Jennifer Schultens, *Tunnel numbers of small knots do not go down under connected sum*, Proc. Amer. Math. Soc. **128** (2000), no. 1, 269–278.
- [NS97] Walter D. Neumann and Gadde A. Swarup, *Canonical decompositions of 3-manifolds*, Geom. Topol. **1** (1997), 21–40 (electronic).
- [Oer84] Ulrich Oertel, *Closed incompressible surfaces in complements of star links*, Pacific J. Math. **111** (1984), no. 1, 209–230.
- [Ota01] Jean-Pierre Otal, *The hyperbolization theorem for fibered 3-manifolds*, SMF/AMS Texts and Monographs, vol. 7, American Mathematical Society, Providence, RI, 2001, Translated from the 1996 French original by Leslie D. Kay.
- [Thu82] William P. Thurston, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.) **6** (1982), no. 3, 357–381.
- [Tsa94] Chichen M. Tsau, *Incompressible surfaces in the knot manifolds of torus knots*, Topology **33** (1994), no. 1, 197–201.
- [Ush04] Akira Ushijima, *Hyperbolic spatial graphs arising from strongly invertible knots*, Topology Appl. **139** (2004), no. 1-3, 253–260.

- [YM84] Shing-Tung Yau and William H. Meeks, III, *The equivariant loop theorem for three-dimensional manifolds and a review of the existence theorems for minimal surfaces*, The Smith conjecture (New York, 1979), Pure Appl. Math., vol. 112, Academic Press, Orlando, FL, 1984, pp. 153–163.